

**LIDOXOR GEL**  
**Topical anesthetic for oral tissues**

**Indications:**

- surface anesthesia of mucosa before injection;
- surface anesthesia in removing primary teeth;
- removing mobile teeth;
- opening periodontal abscess;
- removing dental plaque;

**Contraindications**

Possible allergic reaction to lidocaine.

**Composition and properties**

Sodium carboxymethyl cellulose  
Chamomile extract  
Yarrow extract  
Lidocaine  
Xylitol  
Flavoring agent  
Excipients

The composition of the material includes lidocaine, has a profound and rapid anesthetic effect on the treated surface.

Flavors, contained in the gel, give a pleasant smell, xylitol gives sweet taste. "LIDOXOR" does not cause a burning sensation or tingling. Especially recommended for patients with an allergy to derivatives of of paraaminobenzoic acid, where the risk of such reactions is dramatically reduced.

**Recommended use**

Dry the treated surface of the mucosa. Squeeze out of the tube required amount of gel and smear desired location on the mucosa with the preparation. To achieve a deeper anesthesia it is necessary to apply the gel on a cotton tampon, and then treat with it prepared surface. If necessary, it is possible to apply a tampon to a target surface and keep for some time (1-2 min.) in order to obtain the desired effect.

**Contraindications**

Hypersensitivity to lidocaine and other components of the material.

**Package contents and storage**

Liquid is packed in tubes by 45 ml. "Lidoxor" gel is available in three tastes: green apple, citrus, forest berry. Keep in cool dry place.

Shelf life is 2 years.

After opening use within 1 year.